



## King's Research Portal

DOI:

[10.1016/S2468-2667\(20\)30224-3](https://doi.org/10.1016/S2468-2667(20)30224-3)

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

McCarthy, K., Carter, B., Myint, P., Quinn, T. J., & Hewitt, J. (2020). Frailty and mortality in patients with COVID-19 – Authors' reply. *The Lancet Public Health*, 5(11), E582. [https://doi.org/10.1016/S2468-2667\(20\)30224-3](https://doi.org/10.1016/S2468-2667(20)30224-3)

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

## Authors' reply

We thank all the research teams who have submitted letters in response to our frailty and COVID-19 paper.<sup>1</sup> This study has generated discussion in scientific journals, lay press, and social media. We welcome the opportunity to further discuss our methods and results.

Laurent, Darvall and colleagues, and Pareek and colleagues, all question the use of the Clinical Frailty Scale (CFS) as a tool for decision making during the COVID-19 pandemic. It was not our intention to promote the practice of basing treatment escalation plans on frailty status. In fact, we cautioned against the use of age alone in establishing treatment type.<sup>2</sup> However, in the UK, CFS scoring was recommended at the government level for triaging patients with COVID-19,<sup>3</sup> and so the exploration of the properties of the CFS in this context seemed timely and relevant. Use of frailty assessments in COVID-19 care is not exclusive to the UK; Rockwood and Theou<sup>4</sup> highlighted the need for all health-care professionals working with patients with COVID-19 to familiarise themselves with frailty assessment.

Laurent requests data on C reactive protein (CRP) in light of their work showing no relationship between CRP and mortality in COVID-19.<sup>5</sup> In contrast, we found that for a pre-specified threshold of CRP of 40 mg/L or more on admission, mortality was 31.9% compared with 15.0% if CRP was lower than 40 mg/L (n=1564; adjusted HR=2.61 [95% CI 1.97–3.45]; p<0.0001). Among those who died, median CRP was 115 (IQR: 63–191) compared with 69 (IQR: 29–140) among patients who survived. Our findings concur with those from other studies, suggesting admission CRP can guide the management of COVID-19.<sup>6</sup>

Thus, our data highlight various risk factors for poor outcome in patients admitted with COVID-19. We limited our data collection to factors initially

believed to influence prognosis. Subsequently, other important risk factors have emerged; for example, obesity and ethnicity. All these data improve our understanding of COVID-19 and might improve its management. However, risk factors should not be used in isolation, but rather should inform care discussions with patients and families. With a second wave of COVID-19 in the UK and many European countries, these discussions might once again become a common part of clinical practice.

Since our paper,<sup>1</sup> other studies of frailty and COVID-19 have been reported. Those that made use of a similar prospective, sequential data collection, and follow-up approach have also found associations between frailty and a poor outcome.<sup>7</sup> We acknowledge that there are many unanswered questions around this topic. Our study was an international collaborative effort, and in this spirit, we are happy to work with any future studies that will enhance our understanding of COVID-19 and frailty.

We declare no competing interests.

Crown Copyright © 2020 Published by Elsevier Ltd.  
This is an Open Access article under the CC BY-NC-ND 4.0 license.

**\*Kathryn McCarthy, Jonathan Hewitt, Phyo K Myint, Terry Quinn, Ben Carter**  
kathryn.mccarthy@nbt.nhs.uk

Southmead Hospital, Bristol BS10 5NB, UK (KMC);  
Division of Population Medicine, Cardiff University,  
Cardiff, UK (JH); Institute of Applied Health Sciences,  
University of Aberdeen, Aberdeen, UK (PKM);  
Institute of Cardiovascular and Medical Sciences,  
University of Glasgow, Glasgow, UK (TQ);  
Department of Biostatistics and Health Informatics,  
Institute of Psychiatry, Psychology and Neuroscience,  
King's College London, London, UK (BC)

- 1 Hewitt J, Carter B, Vilches-Moraga A, et al. The effect of frailty on survival in patients with COVID-19 (COPE): a multicentre, European, observational cohort study. *Lancet Public Health* 2020; **5**: e444–51.
- 2 Moug S, Carter B, Myint PK, et al. Decision-making in COVID-19 and frailty. *Geriatrics (Basel)* 2020; **5**: 30.
- 3 National Institute for Health and Care Excellence. COVID-19 rapid guideline: critical care in adults. <https://www.nice.org.uk/guidance/ng159/chapter/1-Admission-to-hospital> (accessed Oct 8, 2020).
- 4 Rockwood K, Theou O. Using the clinical frailty scale in allocating scarce health care resources. *Can Geriatr J* 2020; **23**: 210–15.

- 5 De Smet R, Mellaerts B, Vandewinckele H et al. Frailty and mortality in hospitalized older adults with COVID-19: retrospective observational study. *J Am Med Dir Assoc* 2020; **21**: 928–32.
- 6 Elshazli RM, Toraih EA, Elgaml A, et al. Diagnostic and prognostic value of hematological and immunological markers in COVID-19 infection: a meta-analysis of 6320 patients. *PLoS ONE* 2020; published online Aug 21. <https://doi.org/10.1371/journal.pone.0238160>.
- 7 Aw D, Woodrow L, Ogliari G, Harwood R. Association of frailty with mortality in older inpatients with COVID-19: a cohort study. *Age Ageing* 2020; published online Aug 10. <https://doi.org/10.1093/ageing/afaa184>.

